

Our Mission. Your Success.



Ivermax[®] Pour-On

Ivermectin for Cattle



PRODUCT DESCRIPTION:

Ivermax[®] Pour-On is an effective pour-on formulation that provides treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice and sarcoptic mange mites in cattle.

- Effective against a wide-range of internal and external parasites
- Restricted Drug in California - Use as directed
- Pour-On formulation
- Saves time and money compared to protecting cattle with multiple products
- Provides safe, proven parasite protection at a competitive price
- Easy to use and eliminates an injection site which reduces carcass trim
- A size to fit any livestock operation

INDICATION:

For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle.

Recommended Dose: 1 mL per 22 lb of body weight.

See label for more details on storage, disposal and directions for use.

PRECAUTIONS:

Do not use on lactating dairy cattle.

Close container tightly when not in use. Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced. Do not use when rain is expected to wet cattle within six hours after treatment. This product is for application to skin surface only. Do not give orally or parenterally. Cloudiness in the formulation may occur when Ivermax[®] (ivermectin) Pour-On is stored at temperatures below 32°F. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy. Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g., caked mud or manure. Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, Ivermax[®] (ivermectin) Pour-On is not recommended for use in species other than cattle. Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

SIZES:

- 250 mL
- 1 Liter
- 2.5 Liter
- 5 Liter
- 20 Liter

For More Information:

info@aspenveterinaryresources.com



Ivermax® Pour-On

Ivermectin for Cattle

PACKAGING LABEL:

aspen
VETERINARY RESOURCES, LTD.

IVERMAX®
(ivermectin)
POUR-ON for Cattle
Contains 5 mg ivermectin/mL
PARASITICIDE

Kills: Roundworms (including Brown Stomach Worm), Lungworms, Grubs, Sucking Lice, Biting Lice, Mange Mites, Horn Flies

WARNING! FLAMMABLE!
KEEP AWAY FROM HEAT, SPARKS, OPEN FLAME,
AND OTHER SOURCES OF IGNITION.

WARNING! NOT FOR USE IN HUMANS.
Restricted Drug (California) - Use Only as Directed

ANADA 200-318, Approved by FDA

Manufactured for:
Aspen Veterinary Resources, Ltd.
Liberty, MO 64068, USA
www.aspenveterinaryresources.com

Product No. 11VE030
81VE036 Rev: 12/14

NET CONTENTS:
84.5 fl oz (2.5 Liter)

IVERMAX®
(ivermectin)
POUR-ON for Cattle

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

INDICATIONS: For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle.

See package insert for the complete indications and use directions.

Recommended Dose: 1 mL per 22 lb of body weight.

RESIDUE WARNING: Cattle must not be treated within 48 days of slaughter for human consumption. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

WARNING! NOT FOR USE IN HUMANS.

This product should not be applied to self or others because it may be irritating to human skin and eyes and absorbed through the skin. To minimize accidental skin contact, the user should wear a long-sleeved shirt and rubber gloves. If accidental skin contact occurs, wash immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water and seek medical attention.

Keep this and all drugs out of the reach of children.

WARNING! FLAMMABLE! KEEP AWAY FROM HEAT, SPARKS, OPEN FLAME AND OTHER SOURCES OF IGNITION.

PRECAUTIONS:
Store at 20°C-25°C (68°F-77°F), with the excursions information included on the package insert and protect from light.

Use only in well-ventilated areas or outdoors.

Close container tightly when not in use.

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced. Do not use when rain is expected to wet cattle within six hours after treatment.

This product is for application to skin surface only. Do not give orally or parenterally.

Cloudiness in the formulation may occur when Ivermax® (ivermectin) Pour-On is stored at temperatures below 32°F. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy.

Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g., caked mud or manure.

Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, Ivermax® (ivermectin) Pour-On is not recommended for use in species other than cattle.

Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

Restricted Drug (California) - Use Only as Directed.

Manufactured for:
Aspen Veterinary Resources, Ltd.
Liberty, MO 64068, USA
www.aspenveterinaryresources.com
A1898 12/14
Product No.: 11VE030
81VE036 Rev: 12/14

TAKE TIME OBSERVE LABEL DIRECTIONS

LOT:
EXP:

0 199355 100449 8

Product Code	Product Size	Case
15703301	250mL	12
15893311	1L	12
14563321 or 15998644	2.5L	4
16998644 or 15997844	5L	2
19815157 or 17998644	20L	1